

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ACTOPLUS MET safely and effectively. See full prescribing information for ACTOPLUS MET.

ACTOPLUS MET (pioglitazone and metformin hydrochloride) tablets, for oral use

Initial U.S. Approval: 2005

WARNING: CONGESTIVE HEART FAILURE and LACTIC ACIDOSIS
See full prescribing information for complete boxed warning.

Congestive Heart Failure

- Thiazolidinediones, including pioglitazone, which is a component of ACTOPLUS MET, cause or exacerbate congestive heart failure in some patients. (5.1)
- After initiation of ACTOPLUS MET, and after dose increases, monitor patients carefully for signs and symptoms of heart failure (e.g., excessive, rapid weight gain, dyspnea, and/or edema). If congestive heart failure develops while taking ACTOPLUS MET, consider discontinuation of ACTOPLUS MET or dosage reduction of pioglitazone in ACTOPLUS MET. (5.1)
- ACTOPLUS MET is not recommended in patients with symptomatic heart failure. (5.1)
- Initiation of ACTOPLUS MET in patients with established New York Heart Association (NYHA) Class III or IV heart failure is contraindicated. (4, 5.1)

Lactic Acidosis

- Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms included malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Laboratory abnormalities included elevated blood lactate levels, anion gap acidosis, increased lactate:pyruvate ratio; and metformin plasma levels generally greater than 5 mcg/mL. (5.2)
- Risk factors include renal impairment, concomitant use of certain drugs, age ≥ 65 years old, radiological studies with contrast, surgery and other procedures, hypoxic states, excessive alcohol intake, and hepatic impairment. Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high-risk groups are provided in the Full Prescribing Information. (5.2)
- If lactic acidosis is suspected, discontinue ACTOPLUS MET and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended. (5.2)

RECENT MAJOR CHANGES

Dosage and Administration

Important Dosage and Administration Information (2.1)	06/2024
Recommended Dosage and Administration (2.2)	06/2024
Recommendations for Use in Patients with Renal Impairment (2.3)	06/2024
Recommendations for Congestive Heart Failure (2.4)	06/2024
Coadministration with Strong CYP2C8 Inhibitors (2.5)	06/2024

INDICATIONS AND USAGE

ACTOPLUS MET is a combination of pioglitazone, a thiazolidinedione agonist of peroxisome proliferator receptor gamma, and metformin hydrochloride (HCl), a biguanide, indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. (1)

Limitations of Use:

Not recommended for treatment of type 1 diabetes or diabetic ketoacidosis. (1)

DOSAGE AND ADMINISTRATION

- Obtain liver tests before initiation. If abnormal, use caution when treating with ACTOPLUS MET, investigate the probable cause, treat (if possible), and follow appropriately. (2.1)
- Take orally with meals to reduce gastrointestinal adverse reactions with metformin (2.10)
- Individualize the starting dose based on the patient's current regimen and titrate the dosage gradually, as needed after assessing therapeutic response and tolerability. The maximum recommended total daily dosage is pioglitazone 45 mg and metformin 2,550 mg. (2.2)

- Recommended starting dosage in patients with NYHA Class I or Class II congestive heart failure is 15 mg of pioglitazone and 850 mg of metformin HCl orally once daily. (2.4)
- Prior to initiation, assess renal function with estimated glomerular filtration rate (eGFR). (2.2)
 - Contraindicated in patients with eGFR below 30 mL/min
 - Initiation is not recommended in patients with eGFR between 30 to 45 mL/min
 - Assess risk/benefit of continuing ACTOPLUS MET if eGFR falls below 45 mL/min
 - Discontinue if eGFR falls below 30 mL/min
- Monitor patients for adverse events related to fluid retention after initiation and dose increases. (2.4)
- ACTOPLUS MET may need to be discontinued at time of, or prior to, iodinated contrast imaging procedures. (2.6)

DOSAGE FORMS AND STRENGTHS

Tablets: 15 mg pioglitazone/850 mg metformin HCl. (3)

CONTRAINDICATIONS

- In patients with established New York Heart Association (NYHA) Class III or IV heart failure at the time of ACTOPLUS MET initiation [see Boxed Warning]. (4)
- In patients with severe renal impairment: (eGFR below 30 mL/min). (4)
- In patients with a history of serious hypersensitivity to pioglitazone, metformin HCl, or any of the excipients in ACTOPLUS MET. (4)
- In patients with acute or chronic metabolic acidosis, including diabetic ketoacidosis. (4)

WARNINGS AND PRECAUTIONS

- Congestive heart failure: Fluid retention may occur and can exacerbate or lead to congestive heart failure. Combination use with insulin and use in congestive heart failure NYHA Class I and II may increase risk. Monitor patients for signs and symptoms. (5.1)
- Edema: Dose-related edema may occur. (5.3)
- Hypoglycemia: Consider a lower dose of insulin or insulin secretagogue to reduce risk of hypoglycemia when used in combination with ACTOPLUS MET. (5.4)
- Hepatic effects: Postmarketing reports of hepatic failure, sometimes fatal. Causality cannot be excluded. If liver injury is detected, promptly interrupt ACTOPLUS MET and assess patient for probable cause, then treat cause if possible, to resolution or stabilization. Do not restart ACTOPLUS MET if liver injury is confirmed and no alternate etiology can be found. (5.5)
- Urinary Bladder Tumors: May increase the risk of bladder cancer. Do not use in patients with active bladder cancer. Use caution when using in patients with a prior history of bladder cancer. (5.6)
- Fractures: Increased incidence in female patients. Apply current standards of care for assessing and maintaining bone health. (5.7)
- Macular edema: Postmarketing reports. Recommend regular eye exams in all patients with diabetes according to current standards of care with prompt evaluation for acute visual changes. (5.8)
- Vitamin B₁₂ deficiency: Metformin may lower vitamin B₁₂ levels. Monitor hematologic parameters annually and vitamin B₁₂ at 2 to 3 year intervals and manage any abnormalities. (5.9)

ADVERSE REACTIONS

Most common adverse reactions (>5%) are upper respiratory tract infection, edema, diarrhea, headache and weight gain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Takeda Pharmaceuticals America, Inc. at 1-877-825-3327 and the ACTOPLUS MET website at: www.actoplusmet.com or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Strong CYP2C8 inhibitors (e.g., gemfibrozil): Limit ACTOPLUS MET dose to 15 mg/850 mg daily. (7.1)
- CYP2C8 inducers (e.g., rifampin) may decrease pioglitazone concentrations. (7.2)
- Carbonic anhydrase inhibitors may increase risk of lactic acidosis. Consider more frequent monitoring. (7.3)
- Drugs that reduce metformin clearance (such as ranolazine, vandetanib, dolutegravir, and cimetidine), may increase the accumulation of metformin. Consider the benefits and risks of concomitant use. (7.4)
- Alcohol: Warn patients against excessive alcohol intake. (7.5)

- Use of insulin secretagogues or insulin use may increase the risk for hypoglycemia and may require dose reduction. (7.6)
- Topiramate may decrease pioglitazone concentrations. (7.8)

-----**USE IN SPECIFIC POPULATIONS**-----

- Females and Males of Reproductive Potential: Advise premenopausal females of the potential for an unintended pregnancy. (8.3)
- Pediatrics: Safety and effectiveness have not been established. (8.4)

- Geriatric Use: Assess renal function more frequently. (8.5)
- Hepatic Impairment: Avoid use in patients with hepatic impairment. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 06/2024

FULL PRESCRIBING INFORMATION: CONTENTS***WARNING: CONGESTIVE HEART FAILURE and LACTIC ACIDOSIS****1 INDICATIONS AND USAGE****2 DOSAGE AND ADMINISTRATION**

- 2.1 Important Dosage and Administration Information
- 2.2 Recommended Dosage and Administration
- 2.3 Recommendations for Use in Patients with Renal Impairment
- 2.4 Recommendations for Congestive Heart Failure
- 2.5 Coadministration with Strong CYP2C8 Inhibitors
- 2.6 Discontinuation for Iodinated Contrast Imaging Procedures

3 DOSAGE FORMS AND STRENGTHS**4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

- 5.1 Congestive Heart Failure
- 5.2 Lactic Acidosis
- 5.3 Edema
- 5.4 Hypoglycemia with Concomitant Use with Insulin or Insulin Secretagogues
- 5.5 Hepatic Effects
- 5.6 Urinary Bladder Tumors
- 5.7 Fractures
- 5.8 Macular Edema
- 5.9 Vitamin B₁₂ Levels

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Strong CYP2C8 Inhibitors
- 7.2 CYP2C8 Inducers
- 7.3 Carbonic Anhydrase Inhibitors

7.4 Drugs that Reduce Metformin Clearance

7.5 Alcohol

7.6 Insulin Secretagogues or Insulin

7.7 Drugs Affecting Glycemic Control

7.8 Topiramate

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

10 OVERDOSAGE**11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

14.1 Patients Who Have Inadequate Glycemic Control with Diet and Exercise Alone

14.2 Patients Previously Treated with Metformin

16 HOW SUPPLIED/STORAGE AND HANDLING**17 PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: CONGESTIVE HEART FAILURE and LACTIC ACIDOSIS

Congestive Heart Failure

- Thiazolidinediones, including pioglitazone, which is a component of ACTOPLUS MET, cause or exacerbate congestive heart failure in some patients [see *Warnings and Precautions (5.1)*].
- After initiation of ACTOPLUS MET, and after dose increases, monitor patients carefully for signs and symptoms of heart failure (e.g., excessive, rapid weight gain, dyspnea, and/or edema). If congestive heart failure develops while taking ACTOPLUS MET, consider discontinuation of ACTOPLUS MET or dosage reduction of pioglitazone in ACTOPLUS MET [see *Warnings and Precautions (5.1)*].
- ACTOPLUS MET is not recommended in patients with symptomatic heart failure [see *Warnings and Precautions (5.1)*].
- Initiation of ACTOPLUS MET in patients with established New York Heart Association (NYHA) Class III or IV heart failure is contraindicated [see *Contraindications (4), Warnings and Precautions (5.1)*].

Lactic Acidosis

- Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. The onset of metformin-associated lactic acidosis is often subtle, accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Metformin-associated lactic acidosis was characterized by elevated blood lactate levels (greater than 5 mmol/L), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate:pyruvate ratio; and metformin plasma levels generally greater than 5 mcg/mL [see *Warnings and Precautions (5.2)*].
- Risk factors for metformin-associated lactic acidosis include renal impairment, concomitant use of certain drugs (e.g., carbonic anhydrase inhibitors such as topiramate), age 65 years old or greater, having a radiological study with contrast, surgery and other procedures, hypoxic states (e.g., acute congestive heart failure), excessive alcohol intake, and hepatic impairment.
- Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high-risk groups are provided in the Full Prescribing Information [see *Dosage and Administration (2.2), Contraindications (4), Warnings and Precautions (5.2), Drug Interactions (7), Use in Specific Populations (8.6, 8.7)*].
- If metformin-associated lactic acidosis is suspected, immediately discontinue ACTOPLUS MET and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended [see *Warnings and Precautions (5.2)*].

1 INDICATIONS AND USAGE

ACTOPLUS MET is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

Limitations of Use

ACTOPLUS MET is not recommended to treat type 1 diabetes mellitus or diabetic ketoacidosis.

2 DOSAGE AND ADMINISTRATION**2.1 Important Dosage and Administration Information**

- Obtain liver tests (serum alanine and aspartate aminotransferases, alkaline phosphatase, and total bilirubin) prior to initiating ACTOPLUS MET [see *Warnings and Precautions* (5.5)].
- ACTOPLUS MET contains 15 mg of pioglitazone and 850 mg of metformin hydrochloride (HCl) in each tablet.
- Take ACTOPLUS MET with meals to reduce gastrointestinal adverse reactions with metformin [see *Adverse Reactions* (6.1)].
- If a dose is missed, do not double the next dose.

2.2 Recommended Dosage and AdministrationRecommended Starting Dosage Based on Current Regimen

Individualize the starting dosage of ACTOPLUS MET based on the patient's current regimen and the available strength of ACTOPLUS MET (see Table 1).

Table 1: Recommended Starting Dosage Based on the Patient's Current Regimen

Current Regimen	Starting Dosage of ACTOPLUS MET (15 mg of pioglitazone and 850 mg of metformin HCl per tablet)*
Not treated with either pioglitazone or metformin HCl	One tablet orally once daily
Metformin HCl	One tablet orally once or twice daily. Select a dosage that is as close as possible to the current dosage of metformin HCl
Pioglitazone	One tablet orally once daily
Pioglitazone and metformin HCl	Select a dosage that is as close as possible to the current dosage of pioglitazone and metformin HCl while not exceeding three tablets orally per day.

*For dosage recommendations for patients with renal impairment and/or congestive heart failure, see *Dosage and Administration* (2.3, 2.4)

Dosage Titration for Additional Glycemic Control

Titrate the ACTOPLUS MET dosage gradually, as needed, after assessing therapeutic response and tolerability.

ACTOPLUS MET may be increased to a maximum recommended total daily dosage of three tablets per day (45 mg of pioglitazone and 2,550 mg of metformin HCl). Total daily dosages of 2,550 mg of metformin HCl may be taken in divided doses three times a day to reduce gastrointestinal adverse reactions [see *Adverse Reactions* (6.1)].

2.3 Recommendations for Use in Patients with Renal Impairment

- Assess renal function prior to initiation of ACTOPLUS MET and periodically thereafter [see *Use in Specific Populations* (8.6)].

- ACTOPLUS MET is contraindicated in patients with an estimated glomerular filtration rate (eGFR) below 30 mL/min.
- Initiation of ACTOPLUS MET in patients with an eGFR between 30 to 45 mL/min is not recommended.
- In patients taking ACTOPLUS MET whose eGFR later falls below 45 mL/min, assess the benefit and risk of continuing therapy.
- Discontinue ACTOPLUS MET if the patient's eGFR later falls below 30 mL/min [see *Contraindications (4), Warnings and Precautions (5.2)*].

2.4 Recommendations for Congestive Heart Failure

Starting Dosage in Patients with NYHA Class I or II Congestive Heart Failure

For patients with preexisting NYHA Class I or II congestive heart failure, the recommended starting dosage of ACTOPLUS MET is 15 mg of pioglitazone and 850 mg of metformin [see *Boxed Warning and Warnings and Precautions (5.1)*].

Monitoring for Fluid Retention and Dosage Modifications for Congestive Heart Failure

After initiation of ACTOPLUS MET or with dosage increase, monitor patients carefully for adverse reactions related to fluid retention as has been seen with pioglitazone (e.g., weight gain, edema and signs and symptoms of congestive heart failure).

If congestive heart failure develops while taking ACTOPLUS MET, consider discontinuation of ACTOPLUS MET or dosage reduction of pioglitazone in ACTOPLUS MET [see *Boxed Warning and Warnings and Precautions (5.1)*].

2.5 Coadministration with Strong CYP2C8 Inhibitors

The maximum recommended dosage of ACTOPLUS MET is one tablet (15 mg of pioglitazone and 850 mg of metformin HCl) once daily when used in combination with gemfibrozil or other strong CYP2C8 inhibitors [see *Drug Interactions (7.1), Clinical Pharmacology (12.3)*].

2.6 Discontinuation for Iodinated Contrast Imaging Procedures

Discontinue ACTOPLUS MET at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min; in patients with a history of liver disease, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure; restart ACTOPLUS MET if renal function is stable [see *Warnings and Precautions (5.2)*].

3 DOSAGE FORMS AND STRENGTHS

- 15 mg of pioglitazone and 850 mg of metformin HCl tablets: White to off-white, oblong, film-coated tablets debossed with "4833M" on one side and "15/850" on the other

4 CONTRAINDICATIONS

ACTOPLUS MET is contraindicated in patients with:

- Established NYHA Class III or IV heart failure at the time of ACTOPLUS MET initiation [see *Boxed Warning*].
- Severe renal impairment (eGFR below 30 mL/min) [see *Warnings and Precautions (5.2)*].
- A history of serious hypersensitivity to pioglitazone, metformin HCl, or any of the excipients in ACTOPLUS MET.

- Acute or chronic metabolic acidosis, including diabetic ketoacidosis [see *Warnings and Precautions (5.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Congestive Heart Failure

Pioglitazone, like other thiazolidinediones, can cause dose-related fluid retention when used alone or in combination with other antidiabetic medications and is most common when pioglitazone is used in combination with insulin. Fluid retention may lead to or exacerbate congestive heart failure. Observe patients for signs and symptoms of congestive heart failure. If congestive heart failure develops while taking ACTOPLUS MET, consider discontinuation of ACTOPLUS MET or dosage reduction of pioglitazone in ACTOPLUS MET [see *Boxed Warning, Contraindications (4), Adverse Reactions (6.1)*].

5.2 Lactic Acidosis

Lactic Acidosis

There have been postmarketing cases of metformin-associated lactic acidosis, including fatal cases. These cases had a subtle onset and were accompanied by nonspecific symptoms such as malaise, myalgias, abdominal pain, respiratory distress, or increased somnolence; however, hypothermia, hypotension and resistant bradyarrhythmias have occurred with severe acidosis. Metformin-associated lactic acidosis was characterized by elevated blood lactate concentrations (greater than 5 mmol/L), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate:pyruvate ratio, and metformin plasma levels generally greater than 5 mcg/mL. Metformin decreases liver uptake of lactate increasing lactate blood levels which may increase the risk of lactic acidosis, especially in patients at risk.

If metformin-associated lactic acidosis is suspected, general supportive measures should be instituted promptly in a hospital setting, along with immediate discontinuation of ACTOPLUS MET. In ACTOPLUS MET-treated patients with a diagnosis or strong suspicion of lactic acidosis, prompt hemodialysis is recommended to correct the acidosis and remove accumulated metformin (metformin HCl is dialyzable, with a clearance of up to 170 mL/min under good hemodynamic conditions). Hemodialysis has often resulted in reversal of symptoms and recovery.

Educate patients and their families about the symptoms of lactic acidosis and if these symptoms occur instruct them to discontinue ACTOPLUS MET and report these symptoms to their healthcare provider.

For each of the known and possible risk factors for metformin-associated lactic acidosis, recommendations to reduce the risk of and manage metformin-associated lactic acidosis are provided below:

Renal Impairment

The postmarketing metformin-associated lactic acidosis cases primarily occurred in patients with significant renal impairment. The risk of metformin accumulation and metformin-associated lactic acidosis increases with the severity of renal impairment because metformin is substantially excreted by the kidney. Clinical recommendations based upon the patient's renal function include [see *Dosage and Administration (2.3), Clinical Pharmacology (12.3)*].

- Before initiating ACTOPLUS MET, obtain an eGFR.
- ACTOPLUS MET is contraindicated in patients with an eGFR less than 30 mL/min. Initiation of ACTOPLUS MET is not recommended in patients with eGFR between 30 to 45 mL/min [see *Contraindications (4)*].

- Obtain an eGFR at least annually in all patients taking ACTOPLUS MET. In patients at increased risk for the development of renal impairment (e.g., the elderly), renal function should be assessed more frequently.
- In patients taking ACTOPLUS MET whose eGFR later falls below 45 mL/min, assess the benefit and risk of continuing therapy.

Drug Interactions

The concomitant use of ACTOPLUS MET with specific drugs may increase the risk of metformin-associated lactic acidosis: those that impair renal function, result in significant hemodynamic change, interfere with acid-base balance or increase metformin accumulation (e.g., cationic drugs) [see *Drug Interactions (7)*]. Therefore, consider more frequent monitoring of patients.

Age 65 or Greater

The risk of metformin-associated lactic acidosis increases with the patient's age because elderly patients have a greater likelihood of having hepatic, renal, or cardiac impairment than younger patients. Assess renal function more frequently in elderly patients [see *Use in Specific Populations (8.5)*].

Radiological Studies with Contrast

Administration of intravascular iodinated contrast agents in metformin-treated patients has led to an acute decrease in renal function and the occurrence of lactic acidosis. Stop ACTOPLUS MET at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min; in patients with a history of hepatic impairment, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure, and restart ACTOPLUS MET if renal function is stable.

Surgery and Other Procedures

Withholding of food and fluids during surgical or other procedures may increase the risk for volume depletion, hypotension and renal impairment. ACTOPLUS MET should be temporarily discontinued while patients have restricted food and fluid intake.

Hypoxic States

Several of the postmarketing cases of metformin-associated lactic acidosis occurred in the setting of acute congestive heart failure (particularly when accompanied by hypoperfusion and hypoxemia). Cardiovascular collapse (shock), acute myocardial infarction, sepsis, and other conditions associated with hypoxemia have been associated with lactic acidosis and may also cause prerenal azotemia. When such events occur, discontinue ACTOPLUS MET.

Excessive Alcohol Intake

Alcohol potentiates the effect of metformin on lactate metabolism and this may increase the risk of metformin-associated lactic acidosis. Warn patients against excessive alcohol intake while receiving ACTOPLUS MET.

Hepatic Impairment

Patients with hepatic impairment have developed with cases of metformin-associated lactic acidosis. This may be due to impaired lactate clearance resulting in higher lactate blood levels. Therefore, avoid use of ACTOPLUS MET in patients with clinical or laboratory evidence of hepatic disease.

5.3 Edema

In controlled clinical trials with pioglitazone, edema was reported more frequently in patients treated with pioglitazone than in placebo-treated patients and is dose related [see *Adverse Reactions (6.1)*]. In postmarketing experience, reports of new onset or worsening of edema have been received.

ACTOPLUS MET should be used with caution in patients with edema. Because thiazolidinediones, including pioglitazone, can cause fluid retention, which can exacerbate or lead to congestive heart failure, ACTOPLUS MET should be used with caution in patients at risk for congestive heart failure. Patients treated with ACTOPLUS MET should be monitored for signs and symptoms of congestive heart failure [see *Boxed Warning, Warnings and Precautions (5.1)*].

5.4 Hypoglycemia with Concomitant Use with Insulin or Insulin Secretagogues

Insulin and insulin secretagogues, such as sulfonylureas, are known to cause hypoglycemia. Therefore, a lower dosage of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with ACTOPLUS MET [see *Drug Interactions (7.6, 7.7)*].

5.5 Hepatic Effects

There have been postmarketing reports of fatal and non-fatal hepatic failure in patients taking pioglitazone, although the reports contain insufficient information necessary to establish the probable cause. There has been no evidence of drug-induced hepatotoxicity in the pioglitazone controlled clinical trial database to date [see *Adverse Reactions (6.1)*].

Patients with type 2 diabetes mellitus may have fatty liver disease or cardiac disease with episodic congestive heart failure, both of which may cause liver test abnormalities, and they may also have other forms of liver disease, many of which can be treated or managed. Therefore, obtaining a liver test panel (serum alanine aminotransferase [ALT], aspartate aminotransferase [AST], alkaline phosphatase, and total bilirubin) and assessing the patient is recommended before initiating ACTOPLUS MET therapy. In patients with abnormal liver tests, ACTOPLUS MET should be initiated with caution.

Measure liver tests promptly in patients who report symptoms that may indicate liver injury, including fatigue, anorexia, right upper abdominal discomfort, dark urine or jaundice. In this clinical context, if the patient is found to have clinically significant liver enzyme elevations (serum ALT greater than three times the ULN) and if abnormal liver tests persist or worsen, ACTOPLUS MET should be interrupted and investigation done to establish the probable cause. ACTOPLUS MET should not be restarted in these patients without another explanation for the liver test abnormalities.

5.6 Urinary Bladder Tumors

Tumors were observed in the urinary bladder of male rats in the two year carcinogenicity study [see *Nonclinical Toxicology (13.1)*]. In addition, during the three year PROactive clinical trial, 14 patients out of 2,605 (0.54%) randomized to pioglitazone and 5 out of 2,633 (0.19%) randomized to placebo were diagnosed with bladder cancer. After excluding patients in whom exposure to study drug was less than one year at the time of diagnosis of bladder cancer, there were 6 (0.23%) cases on pioglitazone and 2 (0.08%) cases on placebo. After completion of the trial, a large subset of patients was observed for up to 10 additional years, with little additional exposure to pioglitazone. During the 13 years of both PROactive and observational follow-up, the occurrence of bladder cancer did not differ between patients randomized to pioglitazone or placebo [Hazard Ratio (HR) = 1.00; (95% Confidence Interval (CI) : 0.59, 1.72)].

Findings regarding the risk of bladder cancer in patients exposed to pioglitazone vary among observational studies; some did not find an increased risk of bladder cancer associated with pioglitazone, while others did.

A large prospective 10 year observational cohort study conducted in the United States (U.S) found no statistically significant increase in the risk of bladder cancer in diabetic patients ever exposed to pioglitazone, compared to those never exposed to pioglitazone [HR = 1.06; (95% CI: 0.89, 1.26)].

A retrospective cohort study conducted with data from the United Kingdom found a statistically significant association between ever exposure to pioglitazone and bladder cancer [HR = 1.63; (95% CI: 1.22, 2.19)].

Associations between cumulative dose or cumulative duration of exposure to pioglitazone and bladder cancer were not detected in some studies including the 10 year observational study in the U.S., but were in others. Inconsistent findings and limitations inherent in these and other studies preclude conclusive interpretations of the observational data.

Pioglitazone may be associated with an increase in the risk of urinary bladder tumors. There are insufficient data to determine whether pioglitazone is a tumor promoter for urinary bladder tumors.

Consequently, ACTOPLUS MET should not be used in patients with active bladder cancer and the benefits of glycemic control versus unknown risks for cancer recurrence with ACTOPLUS MET should be considered in patients with a prior history of bladder cancer.

5.7 Fractures

In PROactive (the Prospective Pioglitazone Clinical Trial in Macrovascular Events), 5,238 patients with type 2 diabetes mellitus and a history of macrovascular disease were randomized to pioglitazone (N=2,605), force-titrated up to 45 mg daily or placebo (N=2,633) in addition to standard of care. During a mean follow-up of 34.5 months, the incidence of bone fracture in females was 5.1% (44/870) for pioglitazone versus 2.5% (23/905) for placebo. This difference was noted after the first year of treatment and persisted during the course of the study. The majority of fractures observed in female patients were nonvertebral fractures including lower limb and distal upper limb. No increase in the incidence of fracture was observed in men treated with pioglitazone (1.7%) versus placebo (2.1%). The risk of fracture should be considered in the care of patients, especially female patients, treated with ACTOPLUS MET and attention should be given to assessing and maintaining bone health according to current standards of care.

5.8 Macular Edema

Macular edema has been reported in postmarketing experience in diabetic patients who were taking pioglitazone or another thiazolidinedione. Some patients presented with blurred vision or decreased visual acuity, but others were diagnosed on routine ophthalmologic examination.

Most patients had peripheral edema at the time macular edema was diagnosed. Some patients had improvement in their macular edema after discontinuation of the thiazolidinedione.

Patients with diabetes should have regular eye exams by an ophthalmologist according to current standards of care. Patients with diabetes who report any visual symptoms should be promptly referred to an ophthalmologist, regardless of the patient's underlying medications or other physical findings [see *Adverse Reactions (6.1)*].

5.9 Vitamin B₁₂ Levels

In metformin clinical trials of 29-week duration, a decrease to subnormal levels of previously normal serum vitamin B₁₂ levels was observed in approximately 7% of patients. Such decrease, possibly due to interference with B₁₂ absorption from the B₁₂-intrinsic factor complex, may be associated with anemia but appears to be rapidly reversible with discontinuation of metformin or vitamin B₁₂ supplementation. Certain individuals (those with inadequate vitamin B₁₂ or calcium intake or absorption) appear to be predisposed to developing subnormal vitamin B₁₂ levels. Measure hematologic parameters on an annual basis and vitamin B₁₂ at 2 to 3 year intervals in patients on ACTOPLUS MET and manage any abnormalities [see *Adverse Reactions* (6.1)].

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed elsewhere in the labeling:

- Congestive heart failure [see *Boxed Warning, Warnings and Precautions* (5.1)]
- Lactic acidosis [see *Boxed Warning, Warnings and Precautions* (5.2)]
- Edema [see *Warnings and Precautions* (5.3)]
- Hypoglycemia with Concomitant Use with Insulin or Insulin Secretagogues [see *Warnings and Precautions* (5.4)]
- Hepatic Effects [see *Warnings and Precautions* (5.5)]
- Urinary Bladder Tumors [see *Warnings and Precautions* (5.6)]
- Fractures [see *Warnings and Precautions* (5.7)]
- Macular Edema [see *Warnings and Precautions* (5.8)]
- Vitamin B₁₂ Levels [see *Warnings and Precautions* (5.9)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Pioglitazone

Over 8,500 patients with type 2 diabetes mellitus have been treated with pioglitazone in randomized, double-blind, controlled clinical trials, including 2,605 patients with type 2 diabetes mellitus and macrovascular disease treated with pioglitazone from the PROactive clinical trial. In these trials, over 6,000 patients have been treated with pioglitazone for six months or longer, over 4,500 patients have been treated with pioglitazone for one year or longer, and over 3,000 patients have been treated with pioglitazone for at least two years.

In six pooled 16 to 26 week placebo-controlled monotherapy and 16 to 24 week add-on combination therapy trials, the incidence of withdrawals due to adverse events was 4.5% for patients treated with pioglitazone and 5.8% for comparator-treated patients. The most common adverse events leading to withdrawal were related to inadequate glycemic control, although the incidence of these events was lower (1.5%) with pioglitazone than with placebo (3.0%).

In the PROactive trial, the incidence of withdrawals due to adverse events was 9.0% for patients treated with pioglitazone and 7.7% for placebo-treated patients. Congestive heart failure was the most common serious adverse event leading to withdrawal occurring in 1.3% of patients treated with pioglitazone and 0.6% of patients treated with placebo.

Common Adverse Events: 16 to 26 Week Monotherapy Trials

A summary of the incidence and type of common adverse events reported in three pooled 16 to 26 week placebo-controlled monotherapy trials of pioglitazone is provided in Table 2. Terms that are reported represent those that occurred at an incidence of >5% and more commonly in patients treated with pioglitazone than in patients who received placebo. None of these adverse events were related to the pioglitazone dose.

Table 2: Three Pooled 16 to 26 Week Placebo-Controlled Clinical Trials of Pioglitazone Monotherapy: Adverse Events Reported at an Incidence >5% and More Commonly in Patients Treated with Pioglitazone than in Patients Treated with Placebo

	% of Patients	
	Placebo N=259	Pioglitazone N=606
Upper Respiratory Tract Infection	8.5	13.2
Headache	6.9	9.1
Sinusitis	4.6	6.3
Myalgia	2.7	5.4
Pharyngitis	0.8	5.1

Common Adverse Events: 16 to 24 Week Add-on Combination Therapy Trials

A summary of the overall incidence and types of common adverse events reported in trials of pioglitazone add-on to metformin is provided in Table 3. Terms that are reported represent those that occurred at an incidence of >5% and more commonly with the highest tested dose of pioglitazone.

Table 3: 16 to 24 Week Clinical Trials of Pioglitazone Add-on to Metformin

	16 Week Placebo-Controlled Trial Adverse Events Reported in >5% of Patients and More Commonly in Patients Treated with Pioglitazone + Metformin than in Patients Treated with Placebo + Metformin	
	% of Patients	
	Placebo + Metformin N=160	Pioglitazone 30 mg + Metformin N=168
Edema	2.5	6.0
Headache	1.9	6.0
	24 Week Non-Controlled Double-Blind Trial Adverse Events Reported in >5% of Patients and More Commonly in Patients Treated with Pioglitazone 45 mg + Metformin than in Patients Treated with Pioglitazone 30 mg + Metformin	
	% of Patients	
	Pioglitazone 30 mg + Metformin N=411	Pioglitazone 45 mg + Metformin N=416
Upper Respiratory Tract Infection	12.4	13.5
Edema	5.8	13.9
Headache	5.4	5.8
Weight Increased	2.9	6.7

Note: The preferred terms of edema peripheral, generalized edema, pitting edema, and fluid retention were combined to form the aggregate term of "edema."

Common Adverse Events: 24 Week ACTOPLUS MET Clinical Trial

Table 4 summarizes the incidence and types of adverse reactions reported in a controlled, 24 week double-blind clinical trial of ACTOPLUS MET dosed twice daily in patients with inadequate glycemic control on diet and exercise (N=600).

Table 4: Adverse Events (≥5% for ACTOPLUS MET) Reported by Patients with Inadequate Glycemic Control on Diet and Exercise in a 24 Week Double-Blind Clinical Trial of ACTOPLUS MET Administered Twice Daily

	% of Patients		
	ACTOPLUS MET 15/850 mg Twice Daily N=201	Pioglitazone 15 mg Twice Daily N=190	Metformin 850 mg Twice Daily N=209
Diarrhea	9.0	2.6	15.3
Headache	5.5	2.6	4.8

In this 24 week trial, abdominal pain was reported in 2.0% of patients in the ACTOPLUS MET group, 1.6% in the pioglitazone monotherapy group and 3.3% in the metformin monotherapy group.

Common Adverse Events: PROactive Trial

A summary of the overall incidence and types of common adverse events reported in the PROactive trial is provided in Table 5. Terms that are reported represent those that occurred at an incidence of >5% and more commonly in patients treated with pioglitazone than in patients who received placebo.

Table 5: PROactive Trial: Incidence and Types of Adverse Events Reported in >5% of Patients Treated with Pioglitazone and More Commonly than Placebo

	% of Patients	
	Placebo N=2,633	Pioglitazone N=2,605
Hypoglycemia	18.8	27.3
Edema	15.3	26.7
Cardiac Failure	6.1	8.1
Pain in Extremity	5.7	6.4
Back Pain	5.1	5.5
Chest Pain	5.0	5.1

Mean duration of patient follow-up was 34.5 months.

Congestive Heart Failure

A summary of the incidence of adverse events related to congestive heart failure is provided in Table 6 for the 16 to 24 week add-on to metformin trials. None of the events were fatal.

Table 6: Treatment-Emergent Adverse Events of Congestive Heart Failure (CHF) Patients Treated with Pioglitazone or Placebo Added on to Metformin

	Number (%) of Patients			
	Placebo-Controlled Trial (16 weeks)		Non-Controlled Double-Blind Trial (24 weeks)	
	Placebo + Metformin N=160	Pioglitazone 30 mg + Metformin N=168	Pioglitazone 30 mg + Metformin N=411	Pioglitazone 45 mg + Metformin N=416
At least one congestive heart failure event	0	1 (0.6%)	0	1 (0.2%)
Hospitalized	0	1 (0.6%)	0	1 (0.2%)

Table 7: Treatment-Emergent Adverse Events of Congestive Heart Failure (CHF)

Patients Treated with Pioglitazone or Placebo Added on to a Sulfonylurea					
	Number (%) of Patients				
	Placebo-Controlled Trial (16 weeks)			Non-Controlled Double-Blind Trial (24 weeks)	
	Placebo + Sulfonylurea a N=187	Pioglitazone 15 mg + Sulfonylurea a N=184	Pioglitazone 30 mg + Sulfonylurea a N=189	Pioglitazone 30 mg + Sulfonylurea N=351	Pioglitazone 45 mg + Sulfonylurea N=351
At least one congestive heart failure event	2 (1.1%)	0	0	1 (0.3%)	6 (1.7%)
Hospitalized	2 (1.1%)	0	0	0	2 (0.6%)
Patients Treated with Pioglitazone or Placebo Added on to Insulin					
	Number (%) of Patients				
	Placebo-Controlled Trial (16 weeks)			Non-Controlled Double-Blind Trial (24 weeks)	
	Placebo + Insulin N=187	Pioglitazone 15 mg + Insulin N=191	Pioglitazone 30 mg + Insulin N=188	Pioglitazone 30 mg + Insulin N=345	Pioglitazone 45 mg + Insulin N=345
At least one congestive heart failure event	0	2 (1.0%)	2 (1.1%)	3 (0.9%)	5 (1.4%)
Hospitalized	0	2 (1.0%)	1 (0.5%)	1 (0.3%)	3 (0.9%)
Patients Treated with Pioglitazone or Placebo Added on to Metformin					
	Number (%) of Patients				
	Placebo-Controlled Trial (16 weeks)		Non-Controlled Double-Blind Trial (24 weeks)		
	Placebo + Metformin N=160	Pioglitazone 30 mg + Metformin N=168	Pioglitazone 30 mg + Metformin N=411	Pioglitazone 45 mg + Metformin N=416	
At least one congestive heart failure event	0	1 (0.6%)	0	1 (0.2%)	
Hospitalized	0	1 (0.6%)	0	1 (0.2%)	

Table 8: Treatment-Emergent Adverse Events of Congestive Heart Failure (CHF) in Patients with NYHA Class II or III Congestive Heart Failure Treated with Pioglitazone or Glyburide

	Number (%) of Subjects	
	Pioglitazone N=262	Glyburide N=256
Death due to cardiovascular causes (adjudicated)	5 (1.9%)	6 (2.3%)
Overnight hospitalization for worsening CHF (adjudicated)	26 (9.9%)	12 (4.7%)
Emergency room visit for CHF (adjudicated)	4 (1.5%)	3 (1.2%)
Patients experiencing CHF progression during study	35 (13.4%)	21 (8.2%)

Congestive heart failure events leading to hospitalization that occurred during the PROactive trial are summarized in Table 9.

Table 9: Treatment-Emergent Adverse Events of Congestive Heart Failure (CHF) in PROactive Trial

	Number (%) of Patients	
	Placebo N=2,633	Pioglitazone N=2,605
At least one hospitalized congestive heart failure event	108 (4.1%)	149 (5.7%)
Fatal	22 (0.8%)	25 (1.0%)
Hospitalized, nonfatal	86 (3.3%)	124 (4.7%)

Cardiovascular Safety

In the PROactive trial, 5,238 patients with type 2 diabetes mellitus and a history of macrovascular disease were randomized to pioglitazone (N=2,605), force-titrated up to 45 mg daily or placebo (N=2,633) in addition to standard of care. Almost all patients (95%) were receiving cardiovascular medications (beta blockers, ACE inhibitors, angiotensin II receptor blockers, calcium channel blockers, nitrates, diuretics, aspirin, statins, and fibrates). At baseline, patients had a mean age of 62 years, mean duration of diabetes of 9.5 years, and mean HbA1c of 8.1%. Mean duration of follow-up was 34.5 months.

The primary objective of this trial was to examine the effect of pioglitazone on mortality and macrovascular morbidity in patients with type 2 diabetes mellitus who were at high risk for macrovascular events. The primary efficacy variable was the time to the first occurrence of any event in a cardiovascular composite endpoint that included all-cause mortality, nonfatal myocardial infarction (MI) including silent MI, stroke, acute coronary syndrome, cardiac intervention including coronary artery bypass grafting or percutaneous intervention, major leg amputation above the ankle, and bypass surgery or revascularization in the leg. A total of 514 (19.7%) patients treated with pioglitazone and 572 (21.7%) placebo-treated patients experienced at least one event from the primary composite endpoint (HR = 0.90; 95% CI: 0.80, 1.02; p=0.10).

Although there was no statistically significant difference between pioglitazone and placebo for the three year incidence of a first event within this composite, there was no increase in mortality or in total macrovascular events with pioglitazone. The number of first occurrences and total individual events contributing to the primary composite endpoint is shown in Table 10.

Table 10: PROactive Trial: Number of First and Total Events for Each Component Within the Cardiovascular Composite Endpoint

Cardiovascular Events	Placebo N=2,633		Pioglitazone N=2,605	
	First Events n (%)	Total events n	First Events n (%)	Total events n
Any event	572 (21.7)	900	514 (19.7)	803
All-cause mortality	122 (4.6)	186	110 (4.2)	177
Nonfatal myocardial infarction (MI)	118 (4.5)	157	105 (4.0)	131
Stroke	96 (3.6)	119	76 (2.9)	92
Acute coronary syndrome	63 (2.4)	78	42 (1.6)	65
Cardiac intervention (CABG/PCI)	101 (3.8)	240	101 (3.9)	195
Major leg amputation	15 (0.6)	28	9 (0.3)	28
Leg revascularization	57 (2.2)	92	71 (2.7)	115

CABG = coronary artery bypass grafting; PCI = percutaneous intervention.

Weight Gain

Dose-related weight gain occurs when pioglitazone is used alone or in combination with other antidiabetic medications. The mechanism of weight gain is unclear but probably involves a combination of fluid retention and fat accumulation.

Tables 11, 12, and 13 summarize the changes in body weight with pioglitazone and placebo in the 16 to 26 week randomized, double-blind monotherapy and 16 to 24 week combination add-on therapy trials, the PROactive trial, and the 24 week ACTOPLUS MET trial.

Table 11: Weight Changes (kg) from Baseline During Randomized, Double-Blind Clinical Trials

		Control Group (Placebo)	Pioglitazone 15 mg	Pioglitazone 30 mg	Pioglitazone 45 mg
		Median (25th, 75th percentile)	Median (25th, 75th percentile)	Median (25th, 75th percentile)	Median (25th, 75th percentile)
Monotherapy (16 to 26 weeks)		-1.4 (-2.7, 0.0) N=256	0.9 (-0.5, 3.4) N=79	1.0 (-0.9, 3.4) N=188	2.6 (0.2, 5.4) N=79
Combination Therapy (16 to 24 weeks)	Sulfonylurea	-0.5 (-1.8, 0.7) N=187	2.0 (0.2, 3.2) N=183	3.1 (1.1, 5.4) N=528	4.1 (1.8, 7.3) N=333
	Metformin	-1.4 (-3.2, 0.3) N=160	N/A	0.9 (-1.3, 3.2) N=567	1.8 (-0.9, 5.0) N=407
	Insulin	0.2 (-1.4, 1.4) N=182	2.3 (0.5, 4.3) N=190	3.3 (0.9, 6.3) N=522	4.1 (1.4, 6.8) N=338

Table 12: Median Change in Body Weight in Patients Treated with Pioglitazone vs Patients Treated with Placebo During the Double-Blind Treatment Period in the PROactive Trial

	Placebo	Pioglitazone
	Median (25th, 75th percentile)	Median (25th, 75th percentile)
Change from baseline to final visit (kg)	-0.5 (-3.3, 2.0) N=2,581	+3.6 (0.0, 7.5) N=2,560

Note: Median exposure for both pioglitazone and placebo was 2.7 years.

Table 13: Weight Changes (kg) from Baseline During Double-Blind Clinical Trial with ACTOPLUS MET in Patients with Inadequate Glycemic Control on Diet and Exercise

	ACTOPLUS MET 15/850 mg Twice Daily	Pioglitazone 15 mg Twice Daily	Metformin 850 mg Twice Daily
	Median (25th, 75th percentile)	Median (25th, 75th percentile)	Median (25th, 75th percentile)
Change from baseline to final visit (kg)	1.00 (-1.0, 3.0) N=198	1.35 (-0.7, 4.1) N=178	-1.00 (-2.6, 0.4) N=203

Note: Trial duration of 24 weeks.

Edema

Edema induced from taking pioglitazone is reversible when pioglitazone is discontinued. The edema usually does not require hospitalization unless there is co-existing congestive heart failure.

In the 24 week ACTOPLUS MET trial, edema was reported in 3.0% of patients in the ACTOPLUS MET group, 4.2% in the pioglitazone monotherapy group, and 1.4% in the metformin monotherapy group.

A summary of the frequency and types of edema adverse events occurring in clinical investigations of pioglitazone is provided in Table 14.

Table 14: Adverse Events of Edema in Patients Treated with Pioglitazone

		Number (%) of Patients			
		Placebo	Pioglitazone 15 mg	Pioglitazone 30 mg	Pioglitazone 45 mg
Monotherapy (16 to 26 weeks)		3 (1.2%) N=259	2 (2.5%) N= 81	13 (4.7%) N= 275	11 (6.5%) N=169
Combined Therapy (16 to 24 weeks)	Sulfonylurea	4 (2.1%) N=187	3 (1.6%) N=184	61 (11.3%) N=540	81 (23.1%) N=351
	Metformin	4 (2.5%) N=160	N/A	34 (5.9%) N=579	58 (13.9%) N=416
	Insulin	13 (7.0%) N=187	24 (12.6%) N=191	109 (20.5%) N=533	90 (26.1%) N=345

Note: The preferred terms of edema peripheral, generalized edema, pitting edema, and fluid retention were combined to form the aggregate term of "edema."

Table 15: Adverse Events of Edema in Patients in the PROactive Trial

Number (%) of Patients	
Placebo N=2,633	Pioglitazone N=2,605
419 (15.9%)	712 (27.3%)

Note: The preferred terms of edema peripheral, generalized edema, pitting edema, and fluid retention were combined to form the aggregate term of "edema."

Hepatic Effects

There has been no evidence of pioglitazone-induced hepatotoxicity in the pioglitazone controlled clinical trial database to date. One randomized, double-blind, three year trial comparing pioglitazone to glyburide as add-on to metformin and insulin therapy was specifically designed to evaluate the incidence of serum ALT elevation to greater than three times the upper limit of the reference range, measured every eight weeks for the first 48 weeks of the trial then every 12 weeks thereafter. A total of 3/1,051 (0.3%) patients treated with pioglitazone and 9/1,046 (0.9%) patients treated with glyburide developed ALT values greater than three times the upper limit of the reference range. None of the patients treated with pioglitazone in the pioglitazone controlled clinical trial database to date have had a serum ALT greater than three times the upper limit of the reference range and a corresponding total bilirubin greater than two times the upper limit of the reference range, a combination predictive of the potential for severe drug-induced liver injury.

Hypoglycemia

In the pioglitazone clinical trials, adverse events of hypoglycemia were reported based on clinical judgment of the investigators and did not require confirmation with fingerstick glucose testing.

In the 16 week add-on to sulfonylurea trial, the incidence of reported hypoglycemia was 3.7% with pioglitazone 30 mg and 0.5% with placebo. In the 16 week add-on to insulin trial, the incidence of reported hypoglycemia was 7.9% with pioglitazone 15 mg, 15.4% with pioglitazone 30 mg, and 4.8% with placebo.

The incidence of reported hypoglycemia was higher with pioglitazone 45 mg compared to pioglitazone 30 mg in both the 24 week add-on to sulfonylurea trial (15.7% vs 13.4%) and in the 24-week add-on to insulin trial (47.8% vs 43.5%).

Three patients in these four trials were hospitalized due to hypoglycemia. All three patients were receiving pioglitazone 30 mg (0.9%) in the 24 week add-on to insulin trial. An additional 14 patients reported severe hypoglycemia (defined as causing considerable interference with patient's usual activities) that did not require hospitalization. These patients were receiving pioglitazone 45 mg in combination with sulfonylurea (n=2) or pioglitazone 30 mg or 45 mg in combination with insulin (n=12).

Urinary Bladder Tumors

Tumors were observed in the urinary bladder of male rats in the two year carcinogenicity study [see *Nonclinical Toxicology (13.1)*]. During the three-year PROactive clinical trial, 14 patients out of 2,605 (0.54%) randomized to pioglitazone and 5 out of 2,633 (0.19%) randomized to placebo were diagnosed with bladder cancer. After excluding patients in whom exposure to study drug was less than one year at the time of diagnosis of bladder cancer, there were 6 (0.23%) cases on pioglitazone and 2 (0.08%) cases on placebo. After completion of the trial, a large subset of patients was observed for up to 10 additional years, with little additional exposure to pioglitazone. During the 13 years of both PROactive and observational follow-up, the occurrence of bladder cancer did not differ between patients randomized to pioglitazone or placebo (HR = 1.00; [95% CI: 0.59 1.72]) [see *Warnings and Precautions (5.6)*].

Metformin HCl

In a double-blind clinical study of metformin in patients with type 2 diabetes mellitus, a total of 141 patients received metformin therapy (up to 2,550 mg per day) and 145 patients received placebo. Adverse reactions reported in greater than 5% of the metformin patients, and that were more common in metformin than placebo-treated patients, are listed in Table 16. In this trial, diarrhea led to discontinuation of study medication in 6% of patients treated with metformin.

Table 16: Most Common Adverse Reactions (>5.0%) in a Placebo-Controlled Clinical Study of Metformin Monotherapy*

Adverse Reaction	Metformin Monotherapy (n=141)	Placebo (n=145)
	% of Patients	
Diarrhea	53.2	11.7
Nausea/Vomiting	25.5	8.3
Flatulence	12.1	5.5
Asthenia	9.2	5.5
Indigestion	7.1	4.1
Abdominal Discomfort	6.4	4.8
Headache	5.7	4.8

* Reactions that were more common in metformin than placebo-treated patients.

Laboratory Abnormalities

Pioglitazone

Hematologic Effects

Pioglitazone may cause decreases in hemoglobin and hematocrit. In placebo-controlled monotherapy trials, mean hemoglobin values declined by 2% to 4% in patients treated with pioglitazone compared with a mean change in hemoglobin of -1% to +1% in placebo-treated patients. These changes primarily occurred within the first four to 12 weeks of therapy and remained relatively constant thereafter. These changes may be related to increased plasma volume associated with pioglitazone therapy and are not likely to be associated with any clinically significant hematologic effects.

Creatine Phosphokinase

During protocol-specified measurement of serum creatine phosphokinase (CPK) in pioglitazone clinical trials, an isolated elevation in CPK to greater than 10 times the upper limit of the reference range was noted in nine (0.2%) patients treated with pioglitazone (values of 2,150 to 11,400 IU/L) and in no comparator-treated patients. Six of these nine patients continued to receive pioglitazone, two patients were noted to have the CPK elevation on the last day of dosing, and one patient discontinued pioglitazone due to the elevation. These elevations resolved without any apparent clinical sequelae. The relationship of these events to pioglitazone therapy is unknown.

Metformin

Vitamin B₁₂ Concentrations

In metformin clinical trials of 29-week duration, a decrease to subnormal levels of previously normal serum vitamin B₁₂ levels was observed in approximately 7% of patients.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of pioglitazone and/or metformin. Because these reactions are reported voluntarily from a population of

uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Pioglitazone

Cardiac Disorders: Rapid increases in weight, edema, congestive heart failure with and without previously known heart disease or concomitant insulin administration

Eye Disorders: New onset or worsening diabetic macular edema with decreased visual acuity

Hepatobiliary Disorders: Fatal and nonfatal hepatic failure

Metformin

Hepatobiliary Disorders: Cholestatic, hepatocellular, and mixed hepatocellular liver injury.

7 DRUG INTERACTIONS

7.1 Strong CYP2C8 Inhibitors

An inhibitor of CYP2C8 (e.g., gemfibrozil) significantly increases the exposure (area under the serum concentration-time curve or AUC) and half-life ($t_{1/2}$) of pioglitazone. Therefore, the maximum recommended dosage of ACTOPLUS MET is 15 mg of pioglitazone and 850 mg of metformin HCl once daily if used in combination with gemfibrozil or other strong CYP2C8 inhibitors [see *Dosage and Administration (2.3)*, *Clinical Pharmacology (12.3)*].

7.2 CYP2C8 Inducers

An inducer of CYP2C8 (e.g., rifampin) may significantly decrease the exposure (AUC) of pioglitazone. Therefore, if an inducer of CYP2C8 is started or stopped during treatment with ACTOPLUS MET, changes in diabetes treatment may be needed based on clinical response without exceeding the maximum recommended daily dosage of ACTOPLUS MET (45 mg of pioglitazone and 2,550 mg of metformin HCl) [see *Clinical Pharmacology (12.3)*].

7.3 Carbonic Anhydrase Inhibitors

Topiramate or other carbonic anhydrase inhibitors (e.g., zonisamide, acetazolamide or dichlorphenamide) frequently causes a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with ACTOPLUS MET may increase the risk for lactic acidosis. Consider more frequent monitoring of these patients.

7.4 Drugs that Reduce Metformin Clearance

Concomitant use of drugs that interfere with common renal tubular transport systems involved in the renal elimination of metformin (e.g., organic cationic transporter-2 [OCT2]/multidrug and toxin extrusion [MATE] inhibitors such as ranolazine, vandetanib, dolutegravir, and cimetidine) could increase systemic exposure to metformin and may increase the risk for lactic acidosis [see *Clinical Pharmacology (12.3)*]. Consider the benefits and risks of concomitant use.

7.5 Alcohol

Alcohol is known to potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake while receiving ACTOPLUS MET.

7.6 Insulin Secretagogues or Insulin

Coadministration of ACTOPLUS MET with an insulin secretagogue (e.g., sulfonylurea) or insulin may increase the risk of hypoglycemia. If hypoglycemia occurs in a patient coadministered ACTOPLUS MET and an insulin secretagogue (e.g., sulfonylurea), the dose of the insulin secretagogue should be reduced.

7.7 Drugs Affecting Glycemic Control

Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control. These drugs include the thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blockers, and isoniazid. When such drugs are administered to a patient receiving ACTOPLUS MET, the patient should be closely observed for loss of blood glucose control. When such drugs are withdrawn from a patient receiving ACTOPLUS MET, the patient should be observed closely for hypoglycemia.

7.8 Topiramate

A decrease in the exposure of pioglitazone and its active metabolites were noted with concomitant administration of pioglitazone and topiramate [see *Clinical Pharmacology (12.3)*]. The clinical relevance of this decrease is unknown; however, when ACTOPLUS MET and topiramate are used concomitantly, monitor patients for adequate glycemic control.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited data with ACTOPLUS MET or pioglitazone in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. Published studies with metformin use during pregnancy have not reported a clear association with metformin and major birth defect or miscarriage risk (see *Data*). There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy [see *Clinical Considerations*].

In animal reproduction studies, no adverse developmental effects were observed when pioglitazone was administered to pregnant rats and rabbits during organogenesis at exposures up to 5 and 35 times the 45 mg clinical dose, respectively, based on body surface area. No adverse developmental effects were observed when metformin was administered to pregnant Sprague Dawley rats and rabbits during the period of organogenesis at doses up to 2 to 6 times, respectively, a 2,000 mg clinical dose, based on body surface area (see *Data*).

The estimated background risk of major birth defects is 6 to 10% in women with pre-gestational diabetes with a HbA1c >7 and has been reported to be as high as 20 to 25% in women with a HbA1c >10. The estimated background risk of miscarriage for the indicated population is unknown. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/Fetal Risk

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, still birth and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, still birth, and macrosomia related morbidity.

Data

Human Data

Published data from postmarketing studies have not reported a clear association with metformin and major birth defects, miscarriage, or adverse maternal or fetal outcomes when metformin was used during pregnancy. However, these studies cannot definitely establish the absence of any metformin-associated risk because of methodological limitations, including small sample size and inconsistent comparator groups.

Animal Data

Pioglitazone and Metformin HCl

Animal reproduction studies were not conducted with the combined products in ACTOPLUS MET. The following data are based on studies conducted with the individual components of ACTOPLUS MET.

Pioglitazone

Pioglitazone administered to pregnant rats during organogenesis did not cause adverse developmental effects at a dose of 20 mg/kg (~5 times the 45 mg clinical dose), but delayed parturition and reduced embryo-fetal viability at 40 and 80 mg/kg, or ≥ 9 times the 45 mg clinical dose, by body surface area. In pregnant rabbits administered pioglitazone during organogenesis, no adverse developmental effects were observed at 80 mg/kg (~35 times the 45 mg clinical dose), but reduced embryo-fetal viability at 160 mg/kg, or ~69 times the 45 mg clinical dose, by body surface area. When pregnant rats received pioglitazone during late gestation and lactation, delayed postnatal development, attributed to decreased body weight, occurred in offspring at maternal doses of 10 mg/kg and above or ≥ 2 times the 45 mg clinical dose, by body surface area.

Metformin HCl

Metformin HCl did not cause adverse developmental effects when administered to pregnant Sprague Dawley rats and rabbits up to 600 mg/kg/day during the period of organogenesis. This represents an exposure of about 2 to 6 times a 2,000 mg clinical dose based on body surface area (mg/m^2) for rats and rabbits, respectively.

8.2 Lactation

Risk Summary

There is no information regarding the presence of ACTOPLUS MET or pioglitazone in human milk, the effects on the breastfed infant, or the effects on milk production. Pioglitazone is present in rat milk; however, due to species-specific differences in lactation physiology, animal data may not reliably predict drug levels in human milk. Limited published studies report that metformin is present in human milk (*see Data*). However, there is insufficient information on the effects of metformin on the breastfed infant and no available information on the effects of metformin on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ACTOPLUS MET and any potential adverse effects on the breastfed infant from ACTOPLUS MET or from the underlying maternal condition.

Data

Published clinical lactation studies report that metformin is present in human milk which resulted in infant doses approximately 0.11% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.13 and 1. However, the studies were not designed to definitely establish the risk of use of metformin during lactation because of small sample size and limited adverse event data collected in infants.

8.3 Females and Males of Reproductive Potential

Discuss the potential for unintended pregnancy with premenopausal women as therapy with ACTOPLUS MET, may result in ovulation in some anovulatory women.

8.4 Pediatric Use

Safety and effectiveness of ACTOPLUS MET in pediatric patients have not been established.

ACTOPLUS MET is not recommended for use in pediatric patients based on adverse effects observed in adults, including fluid retention and congestive heart failure, fractures, and urinary bladder tumors [see *Warnings and Precautions (5.1, 5.3, 5.6, 5.7)*].

8.5 Geriatric Use

Pioglitazone

A total of 92 patients (15.2%) treated with pioglitazone in the three pooled 16 to 26 week double-blind, placebo-controlled, monotherapy trials were ≥ 65 years old and two patients (0.3%) were ≥ 75 years old. In the two pooled 16 to 24 week add-on to sulfonylurea trials, 201 patients (18.7%) treated with pioglitazone were ≥ 65 years old and 19 (1.8%) were ≥ 75 years old. In the two pooled 16 to 24 week add-on to metformin trials, 155 patients (15.5%) treated with pioglitazone were ≥ 65 years old and 19 (1.9%) were ≥ 75 years old. In the two pooled 16 to 24 week add-on to insulin trials, 272 patients (25.4%) treated with pioglitazone were ≥ 65 years old and 22 (2.1%) were ≥ 75 years old.

In PROactive Trial, 1,068 patients (41.0%) treated with pioglitazone were ≥ 65 years old and 42 (1.6%) were ≥ 75 years old.

In pharmacokinetic studies with pioglitazone, no significant differences were observed in pharmacokinetic parameters between elderly and younger patients [see *Clinical Pharmacology (12.3)*].

Although clinical experiences have not identified differences in effectiveness and safety between the elderly (≥ 65 years) and younger patients, these conclusions are limited by small sample sizes for patients ≥ 75 years old.

Metformin HCl

Controlled clinical studies of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and young patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of lactic acidosis. Assess renal function more frequently in elderly patients [see *Warnings and Precautions (5.2), Dosage and Administration (2.2)*].

8.6 Renal Impairment

Metformin is substantially excreted by the kidney, and the risk of metformin accumulation and lactic acidosis increases with the degree of renal impairment. ACTOPLUS MET is contraindicated in severe renal impairment, which includes patients with an eGFR below 30 mL/min [see *Dosage and Administration (2.2), Contraindications (4), Warnings and Precautions (5.2), Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

Use of metformin in patients with hepatic impairment has been associated with some cases of lactic acidosis. ACTOPLUS MET is not recommended in patients with hepatic impairment [see *Warnings and Precautions (5.2)*].

10 OVERDOSAGE

Pioglitazone

During controlled clinical trials, one case of overdose with pioglitazone was reported. A male patient took 120 mg per day for four days, then 180 mg per day for seven days. The patient denied any clinical symptoms during this period.

Metformin HCl


Overdose of metformin HCl has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin HCl has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases [see *Warnings and Precautions (5.2)*]. Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful for removal of accumulated metformin from patients in whom metformin overdose is suspected.

In the event of overdose, contact the Poison Help Line (1-800-222-1222) or a medical toxicologist for additional overdose management recommendations. Appropriate supportive treatment should be initiated according to the patient's clinical signs and symptoms.

11 DESCRIPTION

ACTOPLUS MET tablets are a thiazolidinediones and biguanide combination product that contains two oral antidiabetic medications: pioglitazone HCl and metformin HCl.


Pioglitazone [(±)-5-[[4-[2-(5-ethyl-2-pyridinyl) ethoxy]phenyl]methyl]-2,4-] thiazolidinedione monohydrochloride contains one asymmetric carbon, and the compound is synthesized and used as the racemic mixture. The two enantiomers of pioglitazone interconvert *in vivo*. No differences were found in the pharmacologic activity between the two enantiomers. The structural formula is as shown:



pioglitazone HCl

Pioglitazone HCl is an odorless white crystalline powder that has a molecular formula of $C_{19}H_{20}N_2O_3S \cdot HCl$ and a molecular weight of 392.90 daltons. It is soluble in *N,N*-dimethylformamide, slightly soluble in anhydrous ethanol, very slightly soluble in acetone and acetonitrile, practically insoluble in water, and insoluble in ether.

Metformin HCl (*N,N*-dimethylimidodicarbonimidic diamide HCl) is a white crystalline powder with a molecular formula of $C_4H_{11}N_5 \cdot HCl$ and a molecular weight of 165.62. Metformin HCl is freely soluble in water and is practically insoluble in acetone, ether, and chloroform. The pKa of metformin is 12.4. The pH of a 1% aqueous solution of metformin HCl is 6.68. The structural formula is as shown:



metformin HCl

ACTOPLUS MET is available as a tablet for oral administration containing 15 mg pioglitazone (as the base) with 850 mg metformin HCl (15 mg/850 mg) formulated with the following excipients: povidone USP, microcrystalline cellulose NF, croscarmellose sodium NF, magnesium stearate NF, hypromellose 2910 USP, polyethylene glycol 8000 NF, titanium dioxide USP, and talc USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

ACTOPLUS MET combines two antihyperglycemic agents: pioglitazone and metformin.

Pioglitazone

Pioglitazone is a thiazolidinedione that depends on the presence of insulin for its mechanism of action. Pioglitazone decreases insulin resistance in the periphery and in the liver resulting in increased insulin-dependent glucose disposal and decreased hepatic glucose output. Pioglitazone is not an insulin secretagogue. Pioglitazone is an agonist for peroxisome proliferator-activated receptor-gamma (PPAR γ). PPAR receptors are found in tissues important for insulin action such as adipose tissue, skeletal muscle, and liver. Activation of PPAR γ nuclear receptors modulates the transcription of a number of insulin responsive genes involved in the control of glucose and lipid metabolism.

In animal models of diabetes, pioglitazone reduces the hyperglycemia, hyperinsulinemia, and hypertriglyceridemia characteristic of insulin-resistant states such as type 2 diabetes mellitus. The metabolic changes produced by pioglitazone result in increased responsiveness of insulin-dependent tissues and are observed in numerous animal models of insulin resistance.

Because pioglitazone enhances the effects of circulating insulin (by decreasing insulin resistance), it does not lower blood glucose in animal models that lack endogenous insulin.

Metformin HCl

Metformin HCl improves glucose tolerance in patients with type 2 diabetes mellitus, lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Metformin does not produce hypoglycemia in either patients with type 2 diabetes mellitus or healthy subjects [except in specific circumstances, see *Warnings and Precautions (5.4)*] and does not cause hyperinsulinemia. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may actually decrease.

12.2 Pharmacodynamics

Pioglitazone

Clinical studies demonstrate that pioglitazone improves insulin sensitivity in insulin-resistant patients. Pioglitazone enhances cellular responsiveness to insulin, increases insulin-dependent glucose disposal and improves hepatic sensitivity to insulin. In patients with type 2 diabetes mellitus, the decreased insulin resistance produced by pioglitazone results in lower plasma glucose concentrations, lower plasma insulin concentrations, and lower HbA1c values. In controlled clinical trials, pioglitazone had an additive effect on glycemic control when used in combination with a sulfonylurea, metformin, or insulin [see *Clinical Studies (14)*].

Patients with lipid abnormalities were included in clinical trials with pioglitazone. Overall, patients treated with pioglitazone had mean decreases in serum triglycerides, mean increases in HDL cholesterol, and no consistent mean changes in LDL and total cholesterol. There is no conclusive evidence of macrovascular benefit with pioglitazone [see *Adverse Reactions (6.1)*].

In a 26 week, placebo-controlled, dose-ranging monotherapy study, mean serum triglycerides decreased in the 15 mg, 30 mg, and 45 mg pioglitazone dose groups compared to a mean increase in the placebo group. Mean HDL cholesterol increased to a greater extent in patients treated with pioglitazone than in the placebo-treated patients. There were no consistent differences for LDL and total cholesterol in patients treated with pioglitazone compared to placebo (see Table 17).

Table 17: Lipids in a 26 Week Placebo-Controlled Monotherapy Dose-Ranging Study

	Placebo	Pioglitazone 15 mg Once Daily	Pioglitazone 30 mg Once Daily	Pioglitazone 45 mg Once Daily
Triglycerides (mg/dL)	N=79	N=79	N=84	N=77
Baseline (mean)	263	284	261	260
Percent change from baseline (adjusted mean*)	4.8%	-9.0% [†]	-9.6% [†]	-9.3% [†]
HDL Cholesterol (mg/dL)	N=79	N=79	N=83	N=77
Baseline (mean)	42	40	41	41
Percent change from baseline (adjusted mean*)	8.1%	14.1% [†]	12.2%	19.1% [†]
LDL Cholesterol (mg/dL)	N=65	N=63	N=74	N=62
Baseline (mean)	139	132	136	127
Percent change from baseline (adjusted mean*)	4.8%	7.2%	5.2%	6.0%
Total Cholesterol (mg/dL)	N=79	N=79	N=84	N=77
Baseline (mean)	225	220	223	214
Percent change from baseline (adjusted mean*)	4.4%	4.6%	3.3%	6.4%

* Adjusted for baseline, pooled center, and pooled center by treatment interaction.

[†] p<0.05 vs placebo.

In the two other monotherapy studies (16 weeks and 24 weeks) and in combination therapy studies with metformin (16 weeks and 24 weeks), the results were generally consistent with the data above.

12.3 Pharmacokinetics

Absorption

ACTOPLUS MET

In bioequivalence studies of ACTOPLUS MET 15 mg/500 mg and 15 mg/850 mg, the area under the curve (AUC) and maximum concentration (C_{max}) of both the pioglitazone and the metformin component following a single dose of the combination tablet were bioequivalent to ACTOS 15 mg concomitantly administered with metformin HCl immediate release (500 mg or 850 mg, respectively) tablets under fasted conditions in healthy subjects.

Administration of ACTOPLUS MET 15 mg/850 mg with food resulted in no change in overall exposure of pioglitazone. With metformin there was no change in AUC; however, mean peak serum concentration of metformin was decreased by 28% when administered with food. A

delayed time to peak serum concentration was observed for both components (1.9 hours for pioglitazone and 0.8 hours for metformin) under fed conditions. These changes are not likely to be clinically significant.

Pioglitazone

Following once-daily administration of pioglitazone, steady-state serum concentrations of both pioglitazone and its major active metabolites, M-III (keto derivative of pioglitazone) and M-IV (hydroxyl derivative of pioglitazone), are achieved within seven days. At steady state, M-III and M-IV reach serum concentrations equal to or greater than that of pioglitazone. At steady state, in both healthy volunteers and patients with type 2 diabetes mellitus, pioglitazone comprises approximately 30 to 50% of the peak total pioglitazone serum concentrations (pioglitazone plus active metabolites) and 20 to 25% of the total AUC.

C_{max} , AUC, and trough serum concentrations (C_{min}) for pioglitazone and M-III and M-IV, increased proportionally with administered doses of 15 mg and 30 mg per day.

Following oral administration of pioglitazone, T_{max} of pioglitazone was within two hours. Food delays the T_{max} to three to four hours, but does not alter the extent of absorption (AUC).

Metformin HCl

The absolute bioavailability of a 500 mg metformin tablet given under fasting conditions is approximately 50 to 60%. Studies using single oral doses of metformin tablets of 500 mg to 1500 mg, and 850 mg to 2,550 mg, indicate that there is a lack of dose proportionality with increasing doses, which is due to decreased absorption rather than an alteration in elimination. At usual clinical doses and dosing schedules of metformin, steady-state plasma concentrations of metformin are reached within 24 to 48 hours and are generally <1 mcg/mL. During controlled clinical trials, maximum metformin plasma levels did not exceed 5 mcg/mL, even at maximum doses.

Food decreases the rate and extent of metformin absorption, as shown by a 40% lower mean C_{max} , a 25% lower AUC, and a 35 minute prolongation of T_{max} following administration of a single 850 mg tablet of metformin with food, compared to the same tablet strength administered fasting. The clinical relevance of these decreases is unknown.

Distribution

Pioglitazone

The mean apparent volume of distribution (V_d/F) of pioglitazone following single-dose administration is 0.63 ± 0.41 (mean \pm SD) L/kg of body weight. Pioglitazone is extensively protein bound (>99%) in human serum, principally to serum albumin. Pioglitazone also binds to other serum proteins, but with lower affinity. M-III and M-IV are also extensively bound (>98%) to serum albumin.

Metformin HCl

The V_d/F of metformin following single oral doses of 850 mg immediate-release metformin averaged 654 ± 358 L. Metformin is negligibly bound to plasma proteins. Metformin partitions into erythrocytes, most likely as a function of time.

Elimination

Metabolism

Pioglitazone

Pioglitazone is extensively metabolized by hydroxylation and oxidation; the metabolites also partly convert to glucuronide or sulfate conjugates. Metabolites M-III and M-IV are the major circulating active metabolites in humans.

In vitro data demonstrate that multiple CYP isoforms are involved in the metabolism of pioglitazone which include CYP2C8 and, to a lesser degree, CYP3A4 with additional contributions from a variety of other isoforms, including the mainly extrahepatic CYP1A1. *In vivo* study of pioglitazone in combination with gemfibrozil, a strong CYP2C8 inhibitor, showed that pioglitazone is a CYP2C8 substrate [see *Dosage and Administration (2.3)*, *Drug Interactions (7.1)*]. Urinary 6 β -hydroxycortisol/cortisol ratios measured in patients treated with pioglitazone showed that pioglitazone is not a strong CYP3A4 enzyme inducer.

Metformin HCl

Intravenous single-dose studies in healthy subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism (no metabolites have been identified in humans) nor biliary excretion.

Excretion

Pioglitazone

Following oral administration, approximately 15 to 30% of the pioglitazone dose is recovered in the urine. Renal elimination of pioglitazone is negligible and the drug is excreted primarily as metabolites and their conjugates. It is presumed that most of the oral dose is excreted into the bile either unchanged or as metabolites and eliminated in the feces.

The mean serum half-life ($t_{1/2}$) of pioglitazone and its metabolites (M-III and M-IV) range from three to seven hours and 16 to 24 hours, respectively. Pioglitazone has an apparent clearance, CL/F, calculated to be five to seven L/hr.

Metformin HCl

Renal clearance is approximately 3.5 times greater than creatinine clearance (CrCl), which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination $t_{1/2}$ of approximately 6.2 hours. In blood, the elimination $t_{1/2}$ is approximately 17.6 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

Specific Populations

Geriatric Patients

Pioglitazone

In healthy elderly subjects, C_{max} of pioglitazone was not significantly different, but AUC values were approximately 21% higher than those achieved in younger subjects. The mean $t_{1/2}$ of pioglitazone was also prolonged in elderly subjects (about ten hours) as compared to younger subjects (about seven hours). These changes are not considered clinically relevant.

Metformin HCl

Limited data from controlled pharmacokinetic studies of metformin in healthy elderly subjects suggest that total CL/F is decreased, the $t_{1/2}$ is prolonged, and C_{max} is increased, compared to healthy young subjects. From these data, it appears that the change in metformin pharmacokinetics with aging is primarily accounted for by a change in renal function.

Pediatric Patients

Pioglitazone

Safety and efficacy of pioglitazone in pediatric patients have not been established. ACTOPLUS MET is not recommended for use in pediatric patients [see *Use in Specific Populations (8.4)*].

Metformin HCl

After administration of a single oral metformin 500 mg tablet with food, geometric mean metformin C_{max} and AUC differed less than 5% between pediatric type 2 diabetic patients (12

to 16 years of age) and gender- and weight-matched healthy adults (20 to 45 years of age), and all with normal renal function.

Male and Female Patients

Pioglitazone

The mean C_{max} and AUC values of pioglitazone were increased 20 to 60% in females compared to males. In controlled clinical trials, HbA1c decreases from baseline were generally greater for females than for males (average mean difference in HbA1c 0.5%). Because therapy should be individualized for each patient to achieve glycemic control, no dosage adjustment is recommended based on gender alone.

Metformin HCl

Metformin pharmacokinetic parameters did not differ significantly between normal subjects and patients with type 2 diabetes mellitus when analyzed according to gender (males=19, females=16). Similarly, in controlled clinical studies in patients with type 2 diabetes mellitus, the antihyperglycemic effect of metformin was comparable in males and females.

Racial or Ethnic Groups

Pioglitazone

Pharmacokinetic data among various ethnic groups are not available.

Metformin HCl

No studies of metformin pharmacokinetic parameters according to race have been performed. In controlled clinical studies of metformin in patients with type 2 diabetes mellitus, the antihyperglycemic effect was comparable in Whites (n=249), Blacks or African Americans (n=51), and Hispanics or Latinos (n=24).

Patients with Renal Impairment

Pioglitazone

The serum elimination half-life of pioglitazone, M-III and M-IV remains unchanged in patients with moderate (CrCl 30 to 50 mL/min) and severe (CrCl <30 mL/min) renal impairment when compared to subjects with normal renal function. Therefore, no dosage adjustment in patients with renal impairment is required.

Metformin HCl

In patients with decreased renal function, the plasma and blood $t_{1/2}$ of metformin is prolonged and the renal clearance is decreased [see *Dosage and Administration (2.3)*, *Contraindications (4)*, *Warnings and Precautions (5.2)*].

Patients with Hepatic Impairment

Pioglitazone

Compared with healthy controls, subjects with impaired hepatic function (Child-Turcotte-Pugh Grade B/C) have an approximate 45% reduction in pioglitazone and total pioglitazone (pioglitazone, M-III, and M-IV) mean C_{max} but no change in the mean AUC values. Therefore, no dosage adjustment in patients with hepatic impairment is required.

There are postmarketing reports of liver failure with pioglitazone and clinical trials have generally excluded patients with serum ALT >2.5 times the upper limit of the reference range. Use ACTOPLUS MET with caution in patients with liver disease [see *Warnings and Precautions (5.5)*].

Metformin HCl

No pharmacokinetic studies of metformin have been conducted in subjects with hepatic impairment [see *Warnings and Precautions (5.5)*].

Drug Interaction Studies

Specific pharmacokinetic drug interaction studies with ACTOPLUS MET have not been performed, although such studies have been conducted with the individual pioglitazone and metformin components.

*Pioglitazone***Table 18: Effect of Pioglitazone Coadministration on Systemic Exposure of Other Drugs**

Pioglitazone Dosage Regimen (mg)*	Coadministered Drug				
	Name and Dose Regimens	Change in AUC [†]		Change in C _{max} [†]	
45 mg (N = 12)	Warfarin[‡]				
	Daily loading then maintenance doses based PT and INR values Quick's Value = 35 ± 5%	R-Warfarin	↓3%	R-Warfarin	↓2%
		S-Warfarin	↓1%	S-Warfarin	↑1%
45 mg (N = 12)	Digoxin				
	0.200 mg twice daily (loading dose) then 0.250 mg daily (maintenance dose, 7 days)	↑15%		↑17%	
45 mg daily for 21 days (N = 35)	Oral Contraceptive				
	[Ethinyl Estradiol (EE) 0.035 mg plus Norethindrone (NE) 1 mg] for 21 days	EE	↓11%	EE	↓13%
		NE	↑3%	NE	↓7%
45 mg (N = 23)	Fexofenadine				
	60 mg twice daily for 7 days	↑30%		↑37%	
45 mg (N = 14)	Glipizide				
	5 mg daily for 7 days	↓3%		↓8%	
45 mg daily for 8 days (N = 16)	Metformin				
	1,000 mg single dose on Day 8	↓3%		↓5%	
45 mg (N = 21)	Midazolam				
	7.5 mg single dose on Day 15	↓26%		↓26%	
45 mg (N = 24)	Ranitidine				
	150 mg twice daily for 7 days	↑1%		↓1%	
45 mg daily for 4 days (N = 24)	Nifedipine ER				
	30 mg daily for 4 days	↓13%		↓17%	
45 mg (N = 25)	Atorvastatin Calcium				
	80 mg daily for 7 days	↓14%		↓23%	
45 mg (N = 22)	Theophylline				
	400 mg twice daily for 7 days	↑2%		↑5%	

* Daily for 7 days unless otherwise noted.

[†] % change (with/without coadministered drug and no change = 0%); symbols of ↑ and ↓ indicate the exposure increase and decrease, respectively.

[‡] Pioglitazone had no clinically significant effect on prothrombin time.

Table 19: Effect of Coadministered Drugs on Pioglitazone Systemic Exposure

Coadministered Drug and Dosage Regimen	Pioglitazone		
	Dose Regimen (mg)*	Change in AUC [†]	Change in C _{max} [†]
Gemfibrozil 600 mg twice daily for 2 days (N = 12)	15 mg single dose	↑3.2-fold [‡]	↑6%
Ketoconazole 200 mg twice daily for 7 days (N = 28)	45 mg	↑34%	↑14%
Rifampin 600 mg daily for 5 days (N = 10)	30 mg single dose	↓54%	↓5%
Fexofenadine 60 mg twice daily for 7 days (N = 23)	45 mg	↑1%	0%
Ranitidine 150 mg twice daily for 4 days (N = 23)	45 mg	↓13%	↓16%
Nifedipine ER 30 mg daily for 7 days (N = 23)	45 mg	↑5%	↑4%
Atorvastatin Calcium 80 mg daily for 7 days (N = 24)	45 mg	↓24%	↓31%
Theophylline 400 mg twice daily for 7 days (N = 22)	45 mg	↓4%	↓2%
Topiramate 96 mg twice daily for 7 days [§] (N = 26)	30 mg [§]	↓15% [¶]	0%

* Daily for 7 days unless otherwise noted.

[†] Mean ratio (with/without coadministered drug and no change = 1-fold) % change (with/without coadministered drug and no change = 0%); symbols of ↑ and ↓ indicate the exposure increase and decrease, respectively.

[‡] The half-life of pioglitazone increased from 8.3 hours to 22.7 hours in the presence of gemfibrozil [see *Dosage and Administration (2.3), Drug Interactions (7.1)*].

[§] Indicates duration of concomitant administration with highest twice-daily dose of topiramate from Day 14 onwards over the 22 days of study.

[¶] Additional decrease in active metabolites; 60% for M-III and 16% for M-IV.

Metformin HCl

Table 20: Effect of Coadministered Drug on Plasma Metformin Systemic Exposure

Coadministered Drug	Dose of Coadministered Drug*	Dose of Metformin*	Geometric Mean Ratio (ratio with/without coadministered drug) No effect = 1.00	
			AUC [†]	C _{max}
No dosing adjustments required for the following:				
Glyburide	5 mg	500 mg [§]	0.98 [‡]	0.99 [‡]
Furosemide	40 mg	850 mg	1.09 [‡]	1.22 [‡]
Nifedipine	10 mg	850 mg	1.16	1.21
Propranolol	40 mg	850 mg	0.90	0.94
Ibuprofen	400 mg	850 mg	1.05 [‡]	1.07 [‡]
Drugs that are eliminated by renal tubular secretion may increase the accumulation of metformin [see Warnings and Precautions (5), Drug Interactions (7)].				
Cimetidine	400 mg	850 mg	1.40	1.61
Carbonic anhydrase inhibitors may cause metabolic acidosis [see Warnings and Precautions (5), Drug Interactions (7)].				
Topiramate	100 mg [¶]	500 mg [¶]	1.25 [¶]	1.17

* All metformin and coadministered drugs were given as single doses.

[†] AUC = AUC_{0-∞}.

[‡] Ratio of arithmetic means.

[§] Metformin HCl extended-release tablets, 500 mg.

[¶] At steady state with topiramate 100 mg every 12 hours and metformin 500 mg every 12 hours; AUC = AUC_{0-12h}

Table 21: Effect of Metformin on Coadministered Drug Systemic Exposure

Coadministered Drug	Dose of Coadministered Drug*	Dose of Metformin*	Geometric Mean Ratio (ratio with/without coadministered drug) No effect = 1.00	
			AUC [†]	C _{max}
No dosing adjustments required for the following:				
Glyburide	5 mg	500 mg [§]	0.78 [‡]	0.63 [‡]
Furosemide	40 mg	850 mg	0.87 [‡]	0.69 [‡]
Nifedipine	10 mg	850 mg	1.10 [§]	1.08
Propranolol	40 mg	850 mg	1.01 [§]	0.94
Ibuprofen	400 mg	850 mg	0.97 [¶]	1.01 [¶]
Cimetidine	400 mg	850 mg	0.95 [§]	1.01

* All metformin and coadministered drugs were given as single doses.

[†] AUC = AUC_{0-∞}.

[‡] Ratio of arithmetic means, p-value of difference <0.05.

[§] AUC_{0-24hr} reported.

[¶] Ratio of arithmetic means.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

ACTOPLUS MET

No animal studies have been conducted with ACTOPLUS MET. The following data are based on findings in studies performed with pioglitazone or metformin individually.

Pioglitazone

A two year carcinogenicity study was conducted in male and female rats at oral doses up to 63 mg/kg (approximately 14 times the maximum recommended human oral dose of 45 mg based on mg/m²). Drug-induced tumors were not observed in any organ except for the urinary bladder of male rats. Benign and/or malignant transitional cell neoplasms were observed in male rats at 4 mg/kg/day and above (approximately equal to the maximum recommended human oral dose based on mg/m²). Urinary calculi with subsequent irritation and hyperplasia were postulated as the mechanism for bladder tumors observed in male rats. A two year mechanistic study in male rats utilizing dietary acidification to reduce calculi formation was completed in 2009. Dietary acidification decreased but did not abolish the hyperplastic changes in the bladder. The presence of calculi exacerbated the hyperplastic response to pioglitazone but was not considered the primary cause of the hyperplastic changes.

The relevance to humans of the bladder findings in the male rat cannot be excluded.

A two year carcinogenicity study was also conducted in male and female mice at oral doses up to 100 mg/kg/day (approximately 11 times the maximum recommended human oral dose based on mg/m²). No drug-induced tumors were observed in any organ.

Pioglitazone HCl was not mutagenic in a battery of genetic toxicology studies, including the Ames bacterial assay, a mammalian cell forward gene mutation assay (CHO/HPRT and AS52/XPRT), an *in vitro* cytogenetics assay using CHL cells, an unscheduled DNA synthesis assay, and an *in vivo* micronucleus assay.

No adverse effects upon fertility were observed in male and female rats at oral doses up to 40 mg/kg pioglitazone HCl daily prior to and throughout mating and gestation (approximately nine times the maximum recommended human oral dose based on mg/m²).

Metformin HCl

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1500 mg/kg/day, respectively. These doses are both approximately four times a human daily dose of 2000 mg of the metformin component of ACTOPLUS MET based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

There was no evidence of mutagenic potential of metformin in the following *in vitro* tests: Ames test (*S. typhimurium*), gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose of the metformin component of ACTOPLUS MET based on body surface area comparisons.

13.2 Animal Toxicology and/or Pharmacology

Heart enlargement has been observed in mice (100 mg/kg), rats (4 mg/kg and above) and dogs (3 mg/kg) treated orally with pioglitazone HCl (approximately 11, one, and two times the maximum recommended human oral dose for mice, rats, and dogs, respectively, based on mg/m²). In a one year rat study, drug-related early death due to apparent heart dysfunction occurred at an oral dose of 160 mg/kg/day (approximately 35 times the maximum recommended human oral dose based on mg/m²). Heart enlargement was seen in a 13 week study in monkeys at oral doses of 8.9 mg/kg and above (approximately four times the maximum recommended human oral dose based on mg/m²), but not in a 52 week study at oral doses up to 32 mg/kg (approximately 13 times the maximum recommended human oral dose based on mg/m²).

14 CLINICAL STUDIES

14.1 Patients Who Have Inadequate Glycemic Control with Diet and Exercise Alone

In a 24 week, randomized, double-blind clinical trial, 600 patients with type 2 diabetes mellitus inadequately controlled with diet and exercise alone (mean baseline HbA1c 8.7%) were randomized to ACTOPLUS MET 15/850 mg, pioglitazone 15 mg, or metformin 850 mg twice daily. Statistically significant improvements in HbA1c and fasting plasma glucose (FPG) were observed in patients treated with ACTOPLUS MET compared to either pioglitazone or metformin alone (see Table 22).

Table 22: Glycemic Parameters in 24 Week Study of ACTOPLUS MET in Patients with Type 2 Diabetes Mellitus Inadequately Controlled with Diet and Exercise

Parameter	Treatment Group		
	ACTOPLUS MET 15/850 mg Twice Daily	Pioglitazone 15 mg Twice Daily	Metformin 850 mg Twice Daily
HbA1c (%)	N=188	N=162	N=193
Baseline (mean)	8.9	8.7	8.7
Change from Baseline (adjusted mean*)	-1.8	-1.0	-1.0
Difference between ACTOPLUS MET (adjusted mean*) 95% Confidence Interval		0.9 [†] (0.5, 1.2)	0.8 [†] (0.5, 1.2)
% of patients with HbA1c ≤7%	64	47	39
Fasting Plasma Glucose (mg/dL)	N=196	N=176	N=202
Baseline (mean)	177	171	171
Change from Baseline (adjusted mean*)	-40	-22	-25
Difference between ACTOPLUS MET (adjusted mean*) 95% Confidence Interval		18 [†] (8, 28)	15 [†] (6, 25)

* Adjusted for baseline.

[†] p ≤0.05 vs ACTOPLUS MET.

14.2 Patients Previously Treated with Metformin

The efficacy and safety of pioglitazone as add-on to metformin therapy have been established in two clinical studies [see *Clinical Pharmacology (12.3)*].

The two clinical trials testing pioglitazone as add-on to metformin therapy included patients with type 2 diabetes mellitus on any dose of metformin, either alone or in combination with another antidiabetic agent. All other antidiabetic agents were withdrawn at least three weeks prior to starting study treatment.

In the first trial, 328 patients were randomized to receive either 30 mg of pioglitazone or placebo once daily for 16 weeks in addition to their current metformin regimen. Treatment with pioglitazone as add-on to metformin produced statistically significant improvements in HbA1c and FPG at endpoint compared to placebo add-on to metformin (see *Table 23*).

Table 23: Glycemic Parameters in a 16 Week Placebo-Controlled, Add-on to Metformin Trial

	Placebo + Metformin	Pioglitazone 30 mg + Metformin
Total Population		
HbA1c (%)	N=153	N=161
Baseline (mean)	9.8	9.9
Change from baseline (adjusted mean*)	0.2	-0.6
Difference from placebo + metformin (adjusted mean*) 95% Confidence Interval		-0.8 [†] (-1.2, -0.5)
Fasting Plasma Glucose (mg/dL)		
Baseline (mean)	260	254
Change from baseline (adjusted mean*)	-5	-43
Difference from placebo + metformin (adjusted mean*) 95% Confidence Interval		-38 [†] (-49, -26)

* Adjusted for baseline, pooled center, and pooled center by treatment interaction.

[†] p ≤ 0.05 vs placebo + metformin.

In the second trial, 827 patients were randomized to receive either 30 mg or 45 mg of pioglitazone once daily for 24 weeks in addition to their current metformin regimen. The mean reduction from baseline at Week 24 in HbA1c was 0.8% for the 30 mg dose and 1.0% for the 45 mg dose (see *Table 24*). The mean reduction from baseline at Week 24 in FPG was 38 mg/dL for the 30 mg dose and 51 mg/dL for the 45 mg dose.

Table 24: Glycemic Parameters in a 24 Week Add-on to Metformin Study

	Pioglitazone 30 mg + Metformin	Pioglitazone 45 mg + Metformin
Total Population		
HbA1c (%)	N=400	N=398
Baseline (mean)	9.9	9.8
Change from baseline (adjusted mean*)	-0.8	-1.0
Difference from 30 mg daily pioglitazone + metformin (adjusted mean*) (95% CI)		-0.2 (-0.5, 0.1)
Fasting Plasma Glucose (mg/dL)	N=398	N=399
Baseline (mean)	233	232
Change from baseline (adjusted mean*)	-38	-51
Difference from 30 mg daily pioglitazone + metformin (adjusted mean*) (95% CI)		-12 [†] (-21, -4)

95% CI = 95% confidence interval.

* Adjusted for baseline, pooled center, and pooled center by treatment interaction.

[†] p ≤ 0.05 vs. 30 mg daily pioglitazone + metformin.

The therapeutic effect of pioglitazone in combination with metformin was observed in patients regardless of the metformin dose.

16 HOW SUPPLIED/STORAGE AND HANDLING

ACTOPLUS MET is available in 15 mg pioglitazone (as the base)/850 mg metformin HCl tablets as follows:

15 mg/850 mg tablet: white to off-white, oblong, film-coated tablet with “4833M” on one side and “15/850” on the other, available in:

Bottles of 60	NDC 64764-158-60
Bottles of 180	NDC 64764-158-18

Storage

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Keep container tightly closed, and protect from moisture and humidity.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

- **Congestive Heart Failure:** Inform patients of the signs and symptoms of heart failure. Instruct patients who experience an unusually rapid increase in weight or edema, shortness of breath, or other symptoms of heart failure while on ACTOPLUS MET to immediately report these symptoms to their healthcare provider. [see *Warnings and Precautions* (5.1)].

- Lactic Acidosis: Explain to patients the risks of lactic acidosis, its symptoms and conditions that predispose to its development, as noted in the *Warnings and Precautions (5.2)* section. Advise patients to discontinue ACTOPLUS MET immediately and to promptly notify their healthcare professional if unexplained hyperventilation, myalgia, gastrointestinal symptoms, malaise, unusual somnolence, or other nonspecific symptoms occur.

Counsel patients against excessive alcohol intake and inform patients about the importance of regular testing of renal function while receiving ACTOPLUS MET.

Inform patients about the importance of regular testing of renal function and hematologic parameters when receiving treatment with ACTOPLUS MET

Instruct patients to inform their doctor that they are taking ACTOPLUS MET prior to any surgical or radiological procedure, as temporary discontinuation of ACTOPLUS MET may be required until renal function has been confirmed to be normal.

- Edema: Inform patients that ACTOPLUS MET use can lead to new-onset or worsening of edema. Instruct patients to immediately report symptoms of rapid weight increase or worsening edema to their healthcare provider [*see Warnings and Precautions (5.3)*].
- Hypoglycemia with Concomitant Use with Insulin or Insulin Secretagogues: Inform patients that the risk of hypoglycemia is increased when ACTOPLUS MET is used with insulin or insulin secretagogues (such as a sulfonylurea). Educate patients on the signs and symptoms of hypoglycemia [*see Warnings and Precautions (5.4)*].
- Hepatic Effects: Instruct patients to promptly stop taking ACTOPLUS MET and seek immediate medical advice if they experience signs or symptoms of liver injury (e.g., unexplained nausea, vomiting, abdominal pain, fatigue, anorexia, or dark urine) [*see Warnings and Precautions (5.5)*].
- Urinary Bladder Tumors: Advise patients to promptly report any hematuria, dysuria or urinary urgency that develops or increases during treatment as these may be due to bladder cancer [*see Warnings and Precautions (5.6)*].
- Fractures: Inform female patients about the risk of fractures while taking ACTOPLUS MET. Provide them with information on factors that may contribute to fracture risk [*see Warnings and Precautions (5.7)*].
- Macular Edema: Educate patients on the signs and symptoms of macular edema and advise them to seek medical attention from an ophthalmologist if they experience symptoms of macular edema [*see Warnings and Precautions (5.8)*].
- Vitamin B₁₂ Levels: Inform patients about the importance of obtaining regular hematological laboratory monitoring while receiving ACTOPLUS MET [*see Warnings and Precautions (5.9)*].
- Females of Reproductive Age: Inform female patients that treatment with ACTOPLUS MET may result in an unintended pregnancy in some premenopausal anovulatory females due to its effect on ovulation [*see Use in Specific Populations (8.3)*].
- Missed Dosage: Instruct patients if a dose is missed, not to double their next dose.

Distributed by:

Takeda Pharmaceuticals America, Inc.

Lexington, MA 02421

ACTOS and ACTOPLUS MET are trademarks of Takeda Pharmaceutical Company Limited registered with the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc.

All other trademarks are the property of their respective owners.

©2024 Takeda Pharmaceuticals America, Inc.

APM001 R17

FPI-0350